



台灣精神醫學會
Taiwanese Society of Psychiatry

55th 年會暨學術研討會

攜手開創精神醫療新境界

Creating Changes in Mental Health Service
through Partnership

論文摘要集

特別演講

專題演講

目錄

專題討論

工作坊

節目表

壁報展示

倫理品質法規課程

A-02	重鬱症患者接受 ECT 後的極度	莊淑雯	78
A-03	突發性感音神經性聽力障礙伴隨憂鬱症風險	曾志傑	79
A-04	亨丁頓舞蹈症可能患者的自殺風險	黃式州	80
A-05	慢性腎臟病患之腦部白質及腦萎縮變化與認知功能相關研究	葉怡君	81
A-06	使用抗憂鬱藥物治療重度憂鬱症治療難度之臨床構面及參數	劉子瑜	82
A-07	反覆性輕度感染與難治型憂鬱症	鄭佳尚	84
A-08	一位末期腎病患者的蟲爬妄想症：個案報告	鄭勝允	85
A-09	經期的情緒障礙症婦女之催乳素濃度	顏如佑	86
A-10	某醫學中心之精神科照會分析	蘇振翔	87
老年精神醫學			
B-01	苯二氮平類藥物與 Z-藥物使用與老人跌倒住院的相關： 台灣巢式病例对照研究	余昇文	88
B-02	喪偶對老人憂鬱之影響：台灣長期追蹤之研究	周少華	89
B-03	中老年思覺失調症病人慢性腎臟病的盛行率及危險因子	陳奕剛	90
B-04	以 Rivastigmine 改善失智症神經精神症狀中的 Charles Bonnet Syndrome： 一例報告	黃彥勳	92
B-05	從實徵生理訊號量測與臨床評估探討高齡長者於輕度認知障礙到 失智症之認知與情緒狀態病程進展	黃敬偉	93
B-06	治療老年憂鬱之頑固疼痛的 pregabalin 加強療法：個案報告	葉大全	94
司法精神醫學			
C-01	性侵害案件中兒童證詞的可信度評估	王麟祥	96
C-02	以權利基礎觀點審視中國精神衛生法	吳文正	97
社區精神醫學			
D-01	全國首創的社區精神醫療服務模式——以草屯康養院為例	黃聖林	97
D-02	結構化壓力覺察團體在門診精神官能症病患之應用	楊紹民	99
D-03	台灣精神護理之家住民照護品質生活經驗之主觀經驗探討	蘇淑芳	100
個別心理治療			
E-01	立基於精神分析理論之移情官能症、邊緣型人格障礙與自戀型人格障礙之 心理治療行動中的支持、面質與中立	林明雄	101
E-02	一位女性病患接受心理治療後轉化症症狀頻率下降	陳宏茂	102
E-03	一例男性 DSM-5 邊緣型人格障礙症共病物質使用障礙症病人之 支持動力性心理治療案例報告	劉書琦	103
E-04	新手治療師初接心理治療的困境：案例報告	蔡書箴	104

A-02

The Attitude of Major Depression Patients after Electro Convulsive Therapy

重鬱症患者接受 ECT 後的態度

Shu-Wen Chuang¹, Ching-Hua Lin¹, Szu-Mei Hsiao², Mei-Rou Lu¹, Chun-mei Tang¹

莊淑雯¹、林清華¹、蕭恩美²、盧美柔¹、湯春梅¹

¹ Kaohsiung Municipal Kai-Syuan Psychiatric Hospital

² Graduate Institute of Health Care, Meiho University

¹ 高雄市立凱旋醫院 ² 英和科技大學健康照護系

Objective: This study intended to assess the attitude of M.D cases post ECT(Electro Convulsive Therapy) intervention attitude Scale; although ECT was originated in 1983, a number of studies confirm that ECT is a safe and effective treatment, it still causes fear and negative perception of the public, resulting in missing good treatment of suitable cases.

Methods: This study used SPSS 17.0 statistical software to analyze the data. Samples are 118 inpatients with major depression, the attitude in the assessment over four aspects: 1. ECT patients overall experience and satisfaction 2. Cases concern on ECT. 3. Compared to other medical procedures 4. The difference of to ECT post ECT. All questionnaires were due on the day next to the ECT series complete. **Results:** Findings show that ECT help their condition accounted for 100 cases (85%), if relapses asking for ECT treatment accounted for 93 cases (79%). The first select priority is taking drugs accounted for 57 cases (48%). Overall, 96 patients (81.3%) indicates a change of attitude to receive ECT post ECT. **Conclusion:** This study found that many patients realize that ECT is beneficial, satisfied, but compared with other studies the about psychosis cases attitude, of their satisfaction is relative low, The reasons may be the case load of major depression, the overall function is better than psychosis cases, therefore there are significant differences in satisfaction questionnaires.